Dr.Amitabh Kumar, Dr.Avirupa Ghose, Dr.Sudha K Ganesh, Dr.Jyotirmay Biswas Sankara Nethralaya, 147, Mukundapur, E M Bypass, Kolkata - 700099, West Bengal, India Financial interests: None




Background

Posterior scleritis is a commonly misdiagnosed entity due to protean manifestations To study the clinical profile of posterior scleritis in a tertiary eye care centre in Eastern India

> Retrospective observational non-comparative case series 20 eyes of 18 patients **Duration : January 2009 – December 2014** Detailed history and a comprehensive ocular examination

**Patients** and Methods



Comments

- Average Duration of Treatment: 8.6 Months - Average Duration of Followup: 25.6 Months
- Recurrence (Post cessation of treatment) 8 Patients (44.45%) – Average 9.1 months

0.1

Conclusions

(p = 0.007)

0.117

- Female and unilateral preponderance in the 4th decade
- Periocular pain a common presenting complaint

High Intra Ocular

Pressure

Subtle clinical signs with a relatively good visual acuity leads to delayed diagnosis

Cataract

Deranged Liver function

test

- No systemic associations in our cohort
- Favourable outcome on treatment with systemic steroids with or without immunosuppressants
- Recurrences are common mandating long term treatment and followup